

Indications for a second r-process from Halo stars

B. Pfeiffer and K.-L. Kratz, Institut für Kernchemie, Universität Mainz, Germany, HGF VISTARS

All elements heavier than Fe are synthesized by neutron-capture and subsequent β -decay, either by the s- or the r-process. Whereas the s-process is fairly understood, the r-process lacks a common consensus on the astrophysical conditions. Nevertheless, applying the model-independent "waiting-point approximation", the isotopic Solar system r-process abundances could be described by a superposition of components with weighting factors following power laws for $A = 63$ to 240 [1].

The recent discovery of the rare class of ultra-metal-poor (UMP) r-process element enriched Halo stars yielded a major step forward in the understanding of the r-process. These ancient stars were formed from the collaps of interstellar gas clouds which had been "contaminated" with heavy elements by only a few explosive events. The earlier determinations of elemental abundances in UMP stars as CS22892-052 or CS31082-001 seemed to demonstrate a unique pattern of r-process abundances from the very metal-poor, old stars up to the much younger Solar system over the whole range of elements from ^{32}Ge up to ^{92}U .

Extending the range of observed elements (by applying the Hubble Space telescope for UV spectroscopy) and studying more Halo stars spanning a wider range of metallicities, a more diverse picture emerged. The unique abundance pattern is confirmed for elements above the second r-process maximum. But elements in the range ^{32}Ge to ^{50}Sn exhibit a different behavior. They show underabundances of odd-Z elements compared to even-Z ones (see, e.g., [2]).

In order to see, if there is a scatter of the abundance patterns from star to star, in Fig. 1 ratios of elemental abundances normalised to ^{63}Eu are displayed compared to the UMP star CS22892-052 with metallicity $[\text{Fe}/\text{H}]=-3.1$. CS31082-001 with the slightly higher metallicity of -2.9 shows the same pattern for light and heavy elements. In the less metal-poor stars BD+17°3248 and HD122563 with metallicities of -2.1 and -1.7 , respectively, the lower-Z region is overabundant. And it seems that even the elements ^{56}Ba to ^{60}Nd above the second maximum are affected.

The total r-process abundance from just above the Fe-group elements up to the heaviest ones can be obtained by the operation of two processes: 1) the classical "main" process with high neutron densities / entropies acting already in the UMP stars and 2) a new nucleosynthesis process with low neutron densities operating in less metal-poor stars producing low-Z elements. The LEPP (light element primary process) proposed in [6] is a possible candidate. Network calculations for low entropies are promising [7].

References

- [1] K.-L. Kratz et al., Ap. J. 403 (1993) 216.
 [2] C. Sneden et al., ApJ 591 (2003) 936.




Figure 1: Elemental abundances in halo stars compared to CS22892-052. The ratios are normalized at ^{63}Eu . The data are from [2] for CS22892-052, [3] for CS31082-001, [4] for BD+17°3248, and [5] for HD126238.

- [3] V. Hill et al., A & A 387 (2002) 560.
 [4] J.J. Cowan et al., Ap.J. 572 (2002) 861.
 [5] J. Westin et al., Ap.J. 530 (2000) 783.
 [6] F. Montes et al., submitted to Ap.J.
 [7] K. Farouqi et al., in preparation